Form PTO-1449

## INFORMATION DISCLOSURE CITATION IN AN APPLICATION



#3

Docket No. ORT1060

09/434,965

Serial No.

Applicant

Alain T. Luxembourg et al

Filing Date

Group Art Unit

November 5, 1999

1644

## U.S. PATENT DOCUMENTS

Exam'r Ìnit.	Document No.	Date	Name	Class	Sub Class	File Date
ny	5,521,288	28 MAY <b>/9</b> 96	Linsley et al			29 MAR 94
MD	5,595,881	21 JAN <b>9</b> 97	Kendrick et al			9 AUG 94
2	5,827,737	27 OCT <b>/9</b> 98	Peterson et al			24 JIN 96
M	5,731,160	24 MAR/198	Malief et al			26 MAY 92
M	5,583,031	10 DEC <b>[9</b> 6	Stern			2 FEB 92
						,

## FOREIGN PATENT DOCUMENTS

Exam'r Init.	Document No.	Date	Country	Class	Sub Class	Translate (select one) Yes No	
				_		] 🗆	
						] 🗆	
						] 🗆	

OTHER REFERENCES (include author, title, date, pertinent pages, etc.)

40		Agrawal GB, Linderman JJ (1996) Mathematical modeling of helper T lymphocyte/antigen-presenting cell interactions: analysis of methods for modifying antigen processing and presentation. J. Theor. Biol. 182, 487-504.
WO		Ashman RF (1973) Lymphocyte receptor movement induced by sheep erythrocyte binding. J. Immunol. 111, 212-220.
2		Altman JD, Moss PAH, Goulder JPR, Barouch DH, McHeyzer-Williams MG, Bell JI, McMichael AJ, Davis MM (1996) Phenotypic analysis of antigen-specific T lymphocytes. Science 274, 94-96.
MO		Bellone G, Geuna M, Carbone A, Silvestri S, Foa R, Emanuelli G, Matera L (1995) Regulatory action of prolactin on the in vitro growth of CD34 <sup>+</sup> ve human hemopoietic progenitor cells. J. Cell. Physiol. 163, 221-231.
KUD		Borrow P, Oldstone MBA (1997) Lymphocytic choriomeningitis virus. In: "Viral Pathogenesis", Neal Nathanson et al., ed. Lippincott-Raven Publishers, Philadelphia, pp. 593-627.
MO		Brinkman, Volker; Kinzel, Bernd; Kristofic, Colette. TCR-Independent Activation of Human CD4 <sup>+</sup> 45RO <sup>-</sup> T Cells by Anti-CD28 Plus IL-2; Induction of Clonal Expansion and Priming for a Th2 Phenotype. The Journal of Immunology, June 1, 1996, 156:4100-4106.
W		Cai Z, Sprent J (1996) Influence of antigen dose and costimulation on the primary response of CD8 <sup>+</sup> I cells in vitro. J. Exp. Med. 183, 2247-2257.
N		Cai Z, Brunmark A, Jackson MR, Loh D, Peterson PA, Sprent J (1996) Transfected <i>Drosophila</i> cells as a probe for defining the minimal requirements for stimulating unprimed CD8 <sup>+</sup> T cells. Proc. Natl. Acad. Sci. USA 93, 14736-14741.
Res	)	Corr M, Slanetz AE, Boyd LF, Jelonek MT, Khilko S, Al-Ramadi BK, Sang Kim Y, Maher SE, Bothwell ALM, Margulies DH (1994) T cell receptor-MHC class I peptide interactions: affinity, kinetics, and specificity. Science 265, 946-949.

Mariane 2 17/01

DIPE.	Sheet 2 of 3
A 1 200	De Bruijn MLH, Nieland JD, Schumacher TNM, Ploegh HL, Kast WM, Melief CJM (1992) Mechanisms of induction of primary virus- specific čytotoxic T lymphocyte responses. Eur. J. Immunol. 22, 3013-3020.
a me	Dillon SR, Jameson SC, Fink PJ (1994) Vβ5 <sup>+</sup> T cell receptors skew toward OVA + H-2K <sup>b</sup> recognition. J. Immunol. 152, 1790-1801.
EL JAS	Engelhard VH (1994) Structure of peptides associated with MHC class I molecules. Current Opinion Immunol. 6, 13-23.
ADEMIA M.	Gold MR, DeFranco AL (1994) Biochemistry of B lymphocyte activation. Adv. Immunol. 55, 221- 295.
100	Grupp SA, Snow EC, Harmony JAK (1987) The phosphatidylinositol response is an early event in the physiologically relevant activation of antigen-specific B lymphocytes. Cell. Immunol. 109, 181-191.
MO	Hou S, Hyland L, Ryan KW, Portner A, Doherty, PC (1994) Virus-specific CD8 <sup>+</sup> T-cell memory determined by clonal burst size. Nature 369, 652-654.
MO	Irsch J, Irlenbusch S, Radl J, Burrows PD, Cooper MD, Radbruch AH (1994) Switch recombination in normal IgA <sub>1</sub> <sup>+</sup> B lymphocytes. Proc. Natl. Acad. Sci. USA 91, 1323-1327.
140	Jackson MR, Song ES, Yang Y, Peterson PA (1992) Empty and peptide-containing conformers of class I major histocompatibility complex molecules expressed in <i>Drosophila melanogaster</i> cells Proc. Natl. Acad. Sci. USA 89, 12117-12121.
NO	Kane KP, Goldstein SAN, Mescher MF (1988) Class I alloantigen is sufficient for cytolytic T lymphocyte binding and transmembrane signaling. Eur. J. Immunol. 18, 1925-1929.
mO	Kane KP, Mescher MF (1993) Activation of CD8-dependent cytotoxic T lymphocyte adhesion and degranulation by peptide class I antigen complexes. J. Immunol. 150, 4788-4797.
ho	Kato K, Radbruch A (1993) Isolation and characterization of CD34 <sup>+</sup> hematopoietic stem cells from human peripheral blood by high-gradient magnetic cell sorting. Cytometry 14, 384-392.
M	Klavinskis LS, Tishon A, Oldstone MBA (1989) Efficiency and effectiveness of cloned virus-specific cytotoxic T lymphocytes in vivo. J. Immunol. 143, 2013-2016.
NA	Kranz DM, Tonegawa S, Eisen HN (1994) Attachment of an anti-receptor antibody to non-target cells renders them susceptible to lysis by a clone of cytotoxic T lymphocytes. Proc. Natl. Acad. Sci. USA 81, 7922-7926.
1000	Lau LL, Jamieson BD, Somasundaram T, Ahmed R (1994) Cytotoxic T-cell memory without antigen. Nature 369, 648-652.
MP	Luxembourg AT, Cooper NR (1994) Modulation of signaling via the B cell antigen receptor by CD21, the receptor for C3dg and EBV. J. Immunol. 153, 4448-4467.
ny	Matsui K, Jay Boniface J, Reay PA, Schild H, Fazekas de St. Groth B, Davis MM (1991) Low affinity interaction of peptide-MHC complexes with T cell receptors. Science 254, 1788-1791.
na	Mescher MF (1995) Molecular interactions in the activation of effector and precursor cytotoxic T lymphocytes. Immunol. Rev. 146, 177-210.
My	Moore MD, DiScipio RG, Cooper NR, Nemerow GR (1989) Hydrodynamic, electron microscopic, and ligand-binding analysis of the Epstein-Barr virus/C3dg receptor (CR2). J. Biol. Chem. 264, 20576-20582.
nes	Myers CD, Kriz MK, Sullivan TJ, Vitettta ES (1987)) Antigen-induced changes in phospholipid metabolism in antigen-binding B lymphocytes. J. Immunol. 138, 1705-1711.
NA	Nakanishi M, Brian AA, McConnell HM (1983) Binding of cytotoxic T-lymphocytes to supported lipid monolayers containing trypsinized H-2K <sup>k</sup> . Mol. Immunol. 20, 1227-1231.
mo	Noelle RJ, Snow EC (1990) Cognate interactions between helper T cells and B cells. Immunol. Today 11, 361-368.
M	Oehen S, Waldner H, Kündig TM, Hengartner H, Zinkernagel RM (1992) Antivirally protective cytotoxic T cell memory to lymphocytic choriomeningitis virus is governed by persisting antigen. J. Exp. Med. 176, 1273-1281.
	Radbruch A, Recktenwald D (1995) Detection and isolation of rare cells. Curr. Opin. Immunol. 7, 270-273.
MO	Ramensee HG, Friede T, Stevanovic S (1995) MHC ligands and peptide motifs: first listing. Immunogenetics 41, 178-228.
nyo	Sawada K, Krantz SB, Dai CH, Koury ST, Horn ST, Glick AD, Civin CI (1990) Purification of human blood burst-forming units-erythroid and demonstration of the evolution of erythropoietin receptors. J. Cell. Physiol. 142, 219-230.
no	Sha WC, Nelson CA, Newberry RD, Kranz DM, Russell JH, Loh DY (1988) Selective expression of an antigen receptor on CD8-bearing T lymphocytes in transgenic mice. Nature 335, 271-274.
2	Snow EC, Noelle RJ, Uhr JW, Vitetta ES (1983b) Activation of antigen-enriched B cells. II. Role of linked recognition in B cell proliferation to thymus-dependent antigens. J. Immunol. 130, 614-618.
no	Snow EC, Fetherson JD, Zimmer S (1986) Induction of the c-myc protooncogene after antigen binding to hapten-specific B cells. J. Exp. Med. 164, 944-949.
no	Snow EC, Vitetta ES, Uhr JW. Activation of Antigen-Enriched B Cells. The Journal of Immunolog, Vol. 130, No.2, February 1983.
I vwo	Stein P, Dubois P, Greenblatt D, Howard M (1986) Induction of antigen-specific proliferation in affinity-purified small lymphocytes: requirement for BSF-1 by type 2 but not type 1 thymus-independent antigens. J. Immunol. 136, 2080-2089.

2/17/07

<del></del>	
NO ME	Sun S, Cai Z, Langlade-Demoyen P, Brunmark A, Jackson MR, Peterson PA, Sprent J (1996) Dual function of Drosophila cells as APCs for naive CD8 <sup>+</sup> T cells: implications for tumor immunotherapy. Immunity 4, 555-564.
AUG 2 1 2000	wain, Susan L.; Huston, Gail; Tonkonogy, Susan; Weinberg, Andrew. Transforming Growth Factor-β and IL-4 Cause Helper T Cell Fecursors To Develop Into Distinct Effector Helper Cells That Differ in Lymphokine Secretion Pattern and Cell Surface Phenotype. The January of Immunology, Vol. 147, 2991-3000, 1991.
Ewa Day	Sykulev Y, Brunmark A, Jackson M, Cohen RJ, Peterson PA, Eisen HN (1994a) Kinetics and affinity of reactions between an antigen- specific T cell receptor and peptide-MHC complexes. Immunity 1, 15-22.
NO	Sykulev Y, Brunmark A, Tsomides TJ, Kageyama S, Jackson M, Peterson PA, Eisen HN (1994b) High-affinity reactions between antigen- specific T-cell receptors and peptides associated with allogeneic and syngeneic major histocompatibility complex class I proteins. Proc. Natl. Acad. Sci. USA 91, 11487-11491.
No	Tallquist MD, Pease LR (1995) Alloreactive 2C T cells recognize a self peptide in the context of the mutant K <sup>bm3</sup> molecule. J. Immunol. 155, 2419-2426.
100	Tallquist MD, Yun TJ, Pease LR (1996) A single T cell receptor recognizes structurally distinct MHC/peptide complexes with high specificity. J. Exp. Med. 184, 1017-1026.
ND	Ukada K, Wiesmüller KH, Kienle S, Jung G, Walden P (1996) Self-MHC-restricted peptides recognized by an alloreactive T lymphocyte clone. J. Immunol. 157, 670-678.
W	Weber S, Traunecker A, Oliveri F, Gerhard W, Karjalainen K (1992) Specific low-affinity recognition of major histocompatibility complex plus peptide by soluble T-cell receptor. Nature 356, 793-796.
NO	Wilson HA, Greenblatt D, Poenie M, Finkelman FD, Tsien RY (1987) Crosslinkage of B lymphocyte surface immunoglobulin by anti-Ig or antigen induces prolonged oscillation of intracellular ionized calcium. J. Exp. Med. 166, 601-606.
ND	Wunderlich J, Shearer G (1991) Induction and measurement of cytotoxic T lymphocyte activity. In: "Current Protocols in Immunology" (Coligan JE, Kruisbeek AM, Margulies DH, Shevach EM, Strober W, eds.) John Wiley and Sons, New York, pp. 3.11.1 - 3.11.15.
MO	Zhang X, Giangreco L, Broome HE, Dargan CM, Swain SL (1995) Control of CD4 effector fate: transforming growth factor \$1 and interleukin 2 synergize to prevent apoptosis and promote effector expansion. J. Exp. Med. 182, 699-709.

Examiner:	Date Consider	· 2/17/W	/			
Examiner: Initial if citation considered, whether or not citation is in confirmance with MPEP \$609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to the applicant.						